Applicant: Tasuku Honjo et al.

Serial No.: 09/966,880 Filed

a5

: September 28, 2001

Page

Attorney's Docket No.: 06501-088001 / J1-101DP2PCT-US

	Replace the paragraph beginning at page 52, line 18, with the following rewritten
	paragraph:
	FIG. 22 schematically shows a degree of homology between an amino acid sequence of
Λ2	mouse AID protein (SEQ ID NO:2) and that of human AID protein (SEQ ID NO:8). The parts
W	with a closed box are cytidine and deoxycytidylate deaminase zinc-binding region which is an
	AID protein active region
	Replace the paragraph beginning at page 68, line 26, with the following rewritten
	paragraph:
	PCR was conducted using an expression vector prepared by inserting cDNA coding a
~4	full length mouse AID protein, prepared in Example 5 into a plasmid vector pGEX4T1, as a
u'	template with a pair of primers (Primer 170: SEQ ID NO:16 and primer 181: SEQ ID NO:17,
	according to the standard method described in the above.

Replace the paragraph beginning at page 70, line 16, with the following rewritten paragraph:

/-As a result, human AID protein (SEQ ID NO:8) has extremely high homology in amino acid sequences with a mouse AID protein (SEQ ID NO:2) (FIG. 22). Amino acid sequences in Cytidine and deoxycytidilate deaminase zinc-binding region which is an active region in AID protein (both mouse AID and human AID amino acid residues 56 to 94 of SEQ ID NOs:2 and 8, respectively) were completely consistent (conserved) between mouse and human -